

Generalized Casimir invariants of six dimensional solvable real Lie algebras with five dimensional nilradical

Rutwig Campoamor-Stursberg
 Laboratoire de Mathématiques et Applications
 F.S.T Université de Haute Alsace
 F-68093 Mulhouse (France)
 R.Campoamor@uha.fr

Abstract

We finish the determination of the invariants of the coadjoint representation of six dimensional real Lie algebras, by determining a fundamental set of invariants for the 99 isomorphism classes of solvable Lie algebras with five dimensional nilradical. We also give some results on the invariants of solvable Lie algebras in arbitrary dimension whose nilradical has codimension one.

1 Introduction

Invariant functions of the coadjoint orbit of semisimple Lie algebras were determined and physically interpreted by Racah in 1950, although some precedent existed in the literature, starting with the pioneering work of Casimir in 1931 (which justifies that these functions are known as Casimir invariants) [1]. More specifically, the polynomials in the universal enveloping algebra invariant by the adjoint representation of a Lie algebra were studied in the case of classical groups by Pontryagin, and by Yen for the exceptional groups [2]. For semisimple Lie algebras the number of functionally independent invariants is completely determined by the dimension of the Cartan subalgebra. This fact led to the belief that other groups would provide invariants of similar form. But for solvable Lie algebras the results found were of a very different nature. Neither the number of invariants nor their specific structure can be determined generically. This fact is due to the more complicated structure of solvable Lie algebras, which do not underly to a root theory like the classical case. During the last decades, the invariants of solvable Lie algebras have become more important due to their significance in representation theory of Lie algebras, where they allow to label irreducible representations or to split representations [3]. In physics they have also shown their importance, as for example in the construction of completely integrable hamiltonian dynamical systems or the obtention of quantum numbers for symmetry groups of systems. Invariants have been determined only for a small number of solvable and non-semisimple Lie algebras, like the Poincaré algebra and its subalgebras or solvable Lie algebras in dimensions less or equal to six [4, 5]. Indeed the solvable case is a quite important one, mainly because of the Levi decomposition theorem, which shows that any Lie algebra is the semidirect product of a semisimple subalgebra and its radical, which is solvable. Since the invariants of semisimple algebras are perfectly known, the question reduces to the invariants of the radicals

and the action of the Levi subalgebra on the radicals.

Solvable Lie algebras have been classified over the real and complex numbers up to dimension six, and their invariants for algebras having rank $r \leq 4$ were computed by Ndogmo few years ago [6]. However, this work is incomplete, since the author left out the case of six dimensional Lie algebras whose nilradical is of dimension five. The justification was that their structure is very similar to the five dimensional case with four dimensional nilradical [5]. In certain aspects this omission is justified, since the structure of the invariants is related, but it must be kept in mind that odd dimensional Lie algebras must have at least an invariant, while for even dimensional Lie algebras it can happen that only trivial invariants exist, i.e., that any invariant function is necessarily a constant. Exactly this point makes this case interesting and worthy to be studied in detail, and allows to obtain some results which are interesting for the general case, as well as the fact that this classification has not been revised due to the relative inaccessibility of the original reference, up to the discovery of some errors by Turkowski when finishing the classification [7].

In this paper we determine the invariants of all real Lie algebras in dimension six whose nilradical is of dimension five, and we show how from the tables we can infer some results which can be formulated in arbitrary dimension. In particular, we show how to construct parametrized families of solvable Lie algebras whose nilradical has codimension one and which have no non-trivial invariants.

Unless otherwise stated, any Lie algebra $\mathfrak{g}$ considered here is of finite dimension over the field $\mathbb{K} = \mathbb{R}$.

The notation used for the six dimensional Lie algebras with five dimensional nilradical, as well as the nomenclature for the latter, are extracted from the paper by Mubarakzyanov where they were classified [8]. We also note that these lists contain some minor mistakes, which have been corrected here without explicit mention.

2 The coadjoint representation

Let G be a Lie group and $\mathfrak{g}$ the corresponding Lie algebra. The coadjoint representation of G is given by the mapping:

$$ad^* : G \rightarrow GL(\mathfrak{g}^*) : (ad_g^* F)(x) = F(ad_{g^{-1}}x), \quad g \in G, F \in \mathfrak{g}^*, x \in \mathfrak{g}$$

Definition 1 *A function $F \in C^\infty(\mathfrak{g}^*)$ is called a semi-invariant for the coadjoint representation if*

$$F(ad_{g^{-1}}x) = \chi(g) F(x)$$

for any $g \in G$, χ being a character of $\mathfrak{g}$. In particular, if $\chi(g) = 1$, then F is called an invariant of $\mathfrak{g}$.

Thus F is an invariant if and only if F is constant on each coadjoint orbit. Semi-invariants of the coadjoint representation are of extraordinary importance for the analysis of completely integrable Hamiltonian dynamical systems, as pointed out by Arhangelsky in [9], and were used by Trofimov to determine the invariants of Borel subalgebras of simple Lie algebras [10] by means of the symmetries of root systems.

Definition 2 *A maximal set of functionally independent invariants of $\mathfrak{g}$ is called a fundamental set of invariants.*

The usual method to compute the (semi-)invariants of a Lie algebra is making use of the theory of linear partial differential equations [11, 12, 13]. Let $\{X_1, \dots, X_n\}$ be a basis of $\mathfrak{g}$ and let $\{C_{ij}^k\}$ be its structure constants over this basis. We can represent $\mathfrak{g}$ in the space $C^\infty(\mathfrak{g}^*)$ by the differential operators

$$\hat{X}_i = -C_{ij}^k x_k \frac{\partial}{\partial x_j} \quad (1)$$

where $[X_i, X_j] = C_{ij}^k X_k$ ($1 \leq i < j \leq n$). It is not difficult to verify that the operators $\hat{X}_i$ satisfy the brackets $[\hat{X}_i, \hat{X}_j] = C_{ij}^k \hat{X}_k$. The following result gives the procedure to obtain the invariants of the coadjoint representation:

Theorem 1 *A function $F \in C^\infty(\mathfrak{g}^*)$ is an invariant if and only if it is a solution of the following system:*

$$\{\hat{X}_i F = 0, 1 \leq i \leq n\} \quad (2)$$

Proofs can be found in [13, 14, 15, 16]. This theorem reduces the determination of the invariant to a system of linear first-order partial differential equations. The method has been used by many authors to study the invariants, and it is the one we use here. In particular, this allows to determine the classical Casimir operators of any Lie algebra [1, 17]. If $\mathfrak{A}$ denotes the universal enveloping algebra of $\mathfrak{g}$ and $Z(\mathfrak{A})$ its center, then it is well known that the elements in $Z(\mathfrak{A})$ correspond to the Casimir operators [18]. This set indeed coincides with the set of polynomial invariants of ad^* , while the rational invariants correspond to ratios of polynomials contained in $\mathfrak{A}$. However, the system (2) can also have non-rational solutions, which do not have such an interpretation in terms of the enveloping algebra. Such solutions will be called generalized Casimir invariants.

For any given Lie algebra $\mathfrak{g}$, the number of functionally independent invariants of the coadjoint representation can be computed from the brackets. More specifically, let $(C_{ij}^k x_k)$ be the matrix which represents the commutator table over the basis $\{X_1, \dots, X_n\}$, the C_{ij}^k being the structure constants over this basis. The matrix is clearly skew-symmetric, which implies that the rank is necessarily even.

Proposition 1 [19] *The cardinal $\mathcal{N}$ of a fundamental set of invariants of $\mathfrak{g}$ is given by $\dim \mathfrak{g} - \sup \left\{ \text{rank} (C_{ij}^k x_k)_{1 \leq i < j \leq \dim \mathfrak{g}} \right\}$. In particular, $\dim \mathfrak{g} \equiv \mathcal{N} \pmod{2}$.*

Thus the number of functionally independent invariants is deduced from the dimension of $\mathfrak{g}$ and the maximal rank of its commutation table considered as matrix, and has the same parity as the dimension of the algebra. In general, the number of polynomial solutions of (2) will be strictly lower than $\dim \mathfrak{g} - \sup \left\{ \text{rank} (C_{ij}^k x_k)_{1 \leq i < j \leq \dim \mathfrak{g}} \right\}$, and only for particular classes of Lie algebras, such as semi-simple or nilpotent Lie algebras we will obtain an equality [17, 20, 21]. On the other hand, for algebraic Lie algebras (i.e., the Lie algebras of algebraic groups [22]) all invariants can be chosen as rational functions [23]. In recent years, and due to the impossibility of classifying solvable Lie algebras in dimensions $n \geq 7$, the analysis has been restricted to important classes of algebras, such as solvable Lie algebras having abelian or Heisenberg nilradicals [24, 25] or rigid Lie algebras [26, 36].

We illustrate the determination of the invariants by an example:

Example 1 Let $\mathfrak{g}_{6,65}$ be the six dimensional (real) Lie algebra whose brackets are:

$$\begin{aligned} [X_3, X_5] &= X_1, & [X_4, X_5] &= X_2 \\ [X_1, X_6] &= \lambda X_1 + X_2, & [X_2, X_6] &= \lambda X_2 \\ [X_3, X_6] &= (\lambda - \gamma) X_3 + X_4, & [X_4, X_6] &= (\lambda - \gamma) X_4 \\ [X_5, X_6] &= \gamma X_5 \end{aligned}$$

In this case, system (2) has the form

$$\left. \begin{aligned} \widehat{X}_1 F &= -(\lambda x_1 + x_2) \partial_{x_6} F = 0 & \widehat{X}_2 F &= -\lambda x_2 \partial_{x_6} F = 0 \\ \widehat{X}_3 F &= (-x_1 \partial_{x_5} - ((\lambda - \mu) x_3 + x_4) \partial_{x_6}) F = 0 \\ \widehat{X}_4 F &= (-x_2 \partial_{x_5} - (\lambda - \mu) x_4 \partial_{x_6}) F = 0 \\ \widehat{X}_5 F &= (x_1 \partial_{x_3} + x_2 \partial_{x_4} - \gamma x_5 \partial_{x_6}) F = 0 \\ \widehat{X}_6 F &= (\lambda x_1 + x_2) \partial_{x_1} + \lambda x_2 \partial_{x_2} + ((\lambda - \mu) x_3 + x_4) \partial_{x_3} + \\ &\quad + (\lambda - \mu) x_4 \partial_{x_4} + \gamma x_5 \partial_{x_5} = 0 \end{aligned} \right\} \quad (3)$$

The system reduces to the following

$$\left. \begin{aligned} \widehat{X}'_5 F &= (x_1 \partial_{x_3} + x_2 \partial_{x_4}) F = 0 \\ : \widehat{X}'_6 F &= (\lambda x_1 + x_2) \partial_{x_1} + \lambda x_2 \partial_{x_2} + ((\lambda - \mu) x_3 + x_4) \partial_{x_3} + \\ &\quad + (\lambda - \mu) x_4 \partial_{x_4} = 0 \end{aligned} \right\} \quad (4)$$

It is easy to see that $I_1 = x_2 \exp\left(\frac{-\lambda x_1}{x_2}\right)$ is a solution of the system, since it trivially verifies the first equation. Consider the new variables $u = \ln(x_1 x_4 - x_2 x_3)$ and $v = x_1 x_2^{-1}$. Then $\widehat{X}'_5(u) = \widehat{X}'_5(v) = 0$ and $\widehat{X}'_6 u = (2\lambda - \mu)$, $\widehat{X}'_6(v) = 1$. Then the new equation

$$\partial_u F(u, v) + \frac{\widehat{X}'_6(v)}{\widehat{X}'_6(u)} \partial_v F(u, v) = 0 \quad (5)$$

is known to provide a solution of (4) [28]. Since $(2\lambda - \mu)v - u$ is a solution of (5), the system (4) has the solution $I_2 = (x_1 x_4 - x_2 x_3) \exp\left(\frac{-(2\lambda - \mu)x_1}{x_2}\right)$. Clearly I_1 and I_2 are functionally independent. and since $\mathcal{N}(\mathfrak{g}_{6,65}) = 2$, we have found a fundamental set of solutions.

3 Six dimensional Lie algebras

The problem of obtaining criteria to classify solvable Lie algebras was initiated by Mubarakzyanov in 1962 [29], and successfully applied to the five dimensional case [30]. Solvable Lie algebras $\mathfrak{r}$ admit a decomposition $\mathfrak{r} = \mathfrak{n} \oplus \mathfrak{t}$, where $\mathfrak{n}$ is the nilradical and $[\mathfrak{t}, \mathfrak{t}] \subset \mathfrak{n}$, $[\mathfrak{t}, \mathfrak{n}] \subset \mathfrak{n}$. Since the nilradical must satisfy $\dim \mathfrak{n} \geq \frac{1}{2} \dim \mathfrak{r}$ [29], the classification of six dimensional Lie algebras reduces to three cases, namely, solvable Lie algebras having nilradical of dimension four, five or three, the latter being immediate. The case $\dim \mathfrak{n} = 5$ was solved by Mubarakzyanov in [8], while the classification of the case $\dim \mathfrak{n} = 4$, which finishes the classification of six dimensional Lie algebras, was presented by Turkowski in [7]. The invariants of solvable Lie algebras in dimension $n \leq 5$ and of nilpotent Lie algebras in dimension $n \leq 6$ were computed in [5], while the invariants of six dimensional solvable

algebras with four dimensional nilradical were determined in [6]. In this work, the author postulates that the structure of invariants for six dimensional Lie algebras with a five dimensional nilradical is rather similar to that of five dimensional solvable algebras with four dimensional radical. This assertion is right in what concerns the general structure of the invariants, but does not consider the fact that even dimensional Lie algebras are allowed to have no non-trivial invariant, as in fact happens. The analysis of these classes can be generalized to obtain some criteria for the nonexistence of non-trivial invariants in arbitrary dimension. This fact makes the analysis of these invariants interesting, and finishes the determination of the invariants of real or solvable Lie algebras in dimension $n \leq 6$.

The classification of the solvable algebras $\mathfrak{r}$ such that $\dim \frac{\mathfrak{r}}{\mathfrak{n}} = 1$ is based on the method developed in [31], and provides 99 isomorphism classes. These algebras are ordered and labelled respect to the structure of its nilradical. The representatives of these five dimensional nilpotent Lie algebras are taken from [32]. In listing the six dimensional Lie algebras of [8] and its invariants, we will preserve the nomenclature used there for both the nilradicals and the algebras. The following table presents the nonzero brackets for the nilradicals:

Table 1: Nilradicals in dimension five

Name	Brackets
$5\mathfrak{g}_1$:	abelian algebra
$\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$	$[X_2, X_3] = X_1$.
$\mathfrak{g}_{4,1} + \mathfrak{g}_1$	$[X_1, X_5] = X_2, \quad [X_4, X_5] = X_1$.
$\mathfrak{g}_{5,1}$	$[X_3, X_5] = X_1, \quad [X_4, X_5] = X_2$.
$\mathfrak{g}_{5,2}$	$[X_2, X_5] = X_1, \quad [X_3, X_5] = X_2, \quad [X_4, X_5] = X_3$.
$\mathfrak{g}_{5,3}$	$[X_2, X_4] = X_3, \quad [X_2, X_5] = X_1, \quad [X_4, X_5] = X_2$.
$\mathfrak{g}_{5,4}$	$[X_2, X_4] = X_1, \quad [X_3, X_5] = X_1$.
$\mathfrak{g}_{5,5}$	$[X_3, X_4] = X_1, \quad [X_2, X_5] = X_1, \quad [X_3, X_5] = X_2$.
$\mathfrak{g}_{5,6}$	$[X_3, X_4] = X_1, \quad [X_2, X_5] = X_1, \quad [X_3, X_5] = X_2, \quad [X_4, X_5] = X_3$

Remark 1 *The original list of Mubarakzhanov [8] presents some minor misprints, as well as the omission of some algebras. Here we present corrected lists, taking into account the corrections indicated by Turkowski in [33] and the author.*

3.1 Nilradical isomorphic to $5\mathfrak{g}_1$

There are 12 isomorphism classes of solvable Lie algebras having an abelian nilradical, all of them depending on one, two or three parameters, with the exception of $\mathfrak{g}_{6,5}$. The invariants of solvable Lie algebras with abelian nilradical are one of the few cases where a general theorem can be established. Their number is determined by the following result:

Proposition 2 [34] *If $\mathfrak{r}$ is a solvable Lie algebra with abelian nilradical $\mathfrak{n}$, then*

$$\mathcal{N}(\mathfrak{r}) = 2 \dim \mathfrak{n} - \dim \mathfrak{r}$$

Thus all the algebras in tables 1 and 2 admit four invariants. Moreover, system (2) reduces considerably in this case, and the invariants are gained from an unique equation. The fundamental

set of invariants are formed by rational functions whenever the element X_6 acts diagonally [6], and this happens only for $\mathfrak{g}_{6,1}$. The remaining algebras have at least one invariant of exponential form, although some invariants reduce to rational or polynomial functions for specific values of the parameters. In particular, if an element X_i belongs to the center of the Lie algebra, then the polynomial x_i is an invariant, since it is in the center of the universal enveloping algebra.

3.2 Nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$

The case of a nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$ provides the highest number of isomorphism classes, 25 in total [tables 4-6]. It is immediately seen that the nilradical is isomorphic to the direct product of the 3 dimensional Heisenberg Lie algebra with $\mathbb{K}^2$. This type of maximal nilpotent ideal will therefore have some analogy to the other general class of Lie algebras whose invariants are perfectly determined, (the nilradical being isomorphic to the Heisenberg Lie algebra [25]). In the present case, it is immediate from the tables that all Lie algebras $\mathfrak{g}_{6,13}, \dots, \mathfrak{g}_{6,38}$ have exactly 2 invariants, of which at least one can be chosen as a rational function. An ocular inspection of tables 4.-6. shows that the determination of the invariants for the algebras having $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$ as nilradical reduces to solve the unique equation:

$$(C_{16}^k x_k \partial_{x_1} + C_{46}^k x_k \partial_{x_4} + C_{56}^k x_k \partial_{x_5}) F = 0 \quad (6)$$

which shows that the structure of the nilradical merely implies that an invariant does not depend on the variables x_2 and x_3 . From this we will obtain generally a fundamental set of invariants formed by a rational function and an exponential function.

3.3 Nilradical isomorphic to $\mathfrak{g}_{4,1} + \mathfrak{g}_1$

14 isomorphism classes have $\mathfrak{g}_{4,1} + \mathfrak{g}_1$ as nilradical, six of them depending on a continuous parameter [tables 7 and 8]. The nilradical $\mathfrak{g}_{4,1} + \mathfrak{g}_1$ is easily seen to be the direct product of $\mathbb{K}$ and a four dimensional filiform Lie algebra (i.e., a nilpotent Lie algebra with maximal nilpotence index). The system (2) reduces to the following system of two equations:

$$(x_2 \partial_{x_1} + x_1 \partial_{x_4}) F = 0 \quad (C_{16}^k x_k \partial_{x_1} + C_{26}^k x_k \partial_{x_2} + C_{36}^k x_k \partial_{x_3} + C_{46}^k x_k \partial_{x_4}) F = 0 \quad \} \quad (7)$$

This tells that in general we will obtain a fundamental set of invariants formed by an exponential and a rational function, depending on the action of X_6 on the nilradical. For concrete values of the parameters the exponential function can reduce to a rational, even to a polynomial invariant.

We can deduce a general result starting from this case: let $\mathfrak{g}_{2m}$ be the $2m$ -dimensional nilpotent Lie algebra given by the brackets $[X_1, X_i] = X_{i+1}, 2 \leq i \leq 2m-1$ over the basis $\{X_1, \dots, X_{2m}\}$. Consider a $(2m+2)$ -dimensional solvable Lie algebra $\mathfrak{r}_{2m+2}$ whose nilradical is isomorphic to $\mathfrak{g}_{2m} \oplus \mathbb{K}$. This generalizes in natural manner the Lie algebras $\mathfrak{g}_{6,53}$ to $\mathfrak{g}_{6,70}$.

Theorem 2 *For $m \geq 2$ the number of invariants of $\mathfrak{r}_{2m+2}$ is given by $\mathcal{N}(\mathfrak{r}_{2m+2}) = 2m - 2$. Moreover, whenever $[X_{2m}, X_{2m+2}] \neq 0$ or $[X_{2m+1}, X_{2m+2}] \neq 0$, an invariant F is a solution of the following system:*

$$\left(\sum_{j=2}^{2m-1} x_{j+1} \partial_{x_j} \right) F = 0 \quad \left(\sum_{i,j=2}^{2m+1} C_{i,2m+2}^k x_k \partial_{x_i} \right) F = 0 \quad \} \quad (8)$$

Proof. The number of invariants follows at once from proposition 1, since the matrix $(C_{ij}^k x_k)_{1 \leq i < j \leq 2m+2}$ representing the commutator table has rank four for any m . For the second assertion, observe that since X_{2m} and X_{2m+1} belong to the center of the nilradical, the associated differential operators are

$$\hat{X}_{2m} = -C_{2m,2m+2}^k x_k \partial_{x_{2m+2}} \quad \hat{X}_{2m+1} = -C_{2m+1,2m+2}^k x_k \partial_{x_{2m+2}} \quad \} \quad (9)$$

and if either $[X_{2m}, X_{2m+2}] \neq 0$ or $[X_{2m+1}, X_{2m+2}] \neq 0$, this implies that $\partial_{x_{2m+2}} F = 0$ for any invariant. On the other hand, from $\hat{X}_{2m-1} = x_{2m} \partial_{x_1} - C_{2m-1,2m+2}^k x_k \partial_{x_{2m+2}}$ it also follows that $\partial_{x_1} F = 0$. This reduces system (2) to the previous form. ■

The practical procedure to solve systems like (5) is to solve the first equation and then search for solutions which are also common to the second equations [35].

3.4 Nilradical isomorphic to $\mathfrak{g}_{5,1}$

There are 18 isomorphism classes of six dimensional Lie algebras with nilradical isomorphic to $\mathfrak{g}_{5,1}$ [tables 9 and 10]. From these, only the algebras $\mathfrak{g}_{6,53}$, $\mathfrak{g}_{6,66}$ and $\mathfrak{g}_{6,69}$ do not depend on parameters. $\mathfrak{g}_{6,53}$ is also the only algebra which admits a fundamental set of invariants formed by two polynomials. For the remaining cases, such a set is either formed by a rational and an exponential function or two exponential functions.

3.5 Nilradical isomorphic to $\mathfrak{g}_{5,2}$

There are five isomorphism classes with $\mathfrak{g}_{5,2}$ as nilradical, from which only one, $\mathfrak{g}_{6,71}$, depends on a continuous parameter, while the algebra $\mathfrak{g}_{6,73}$ depends on a parameter having the values ± 1 . It follows immediately from the brackets listed in table 11. that the invariants of these algebras do not depend on the variables x_5 and x_6 , and that any of these algebras satisfies $\mathcal{N} = 2$. With the exception of the algebras $\mathfrak{g}_{6,73}$ and $\mathfrak{g}_{6,75}$, the action of X_6 on the nilradical is diagonal, which implies that we can always find a fundamental set of invariants formed by rational functions.

3.6 Nilradical isomorphic to $\mathfrak{g}_{5,3}$

There are 6 isomorphism classes having $\mathfrak{g}_{5,3}$ as nilradical, one of them depending on a continuous parameter. It follows immediately from the structure of these algebras that there are exactly two functionally independent invariants. Only for $\mathfrak{g}_{76}$ and $\mathfrak{g}_{6,80}$ the action of X_6 is diagonal, thus only for these two algebras we obtain a fundamental set of invariants formed by rational functions. It is immediately seen from table 12. that system (2) reduces to the following:

$$x_3 \partial_{x_4} + x_1 \partial_{x_5} = 0 \quad x_3 \partial_{x_2} - x_2 \partial_{x_5} = 0 \quad x_1 \partial_{x_2} + x_2 \partial_{x_4} = 0 \quad \sum_{i=1}^5 C_{i6}^k x_k \partial_{x_i} = 0 \quad \} \quad (10)$$

From this we deduce that any invariant can be represented as a function $F\left(x_1, x_3, \frac{2(x_3 x_5 - x_1 x_4) + x_2^2}{x_3}\right)$, the new variables being a fundamental set of solutions for the three first equations, since in any case $\frac{\partial F}{\partial x_6} = 0$. Even more is true: we can choose one of the fundamental invariants as a solution of the equation

$$a x_1 \partial_{x_1} + (b x_1 + c x_3) \partial_{x_3} = 0 \quad (11)$$

depending only on x_1 and x_3 , where a, b, c are constants. Up to the case where $a = c$ and $b = 1$, which provides an exponential solution, the remaining cases have the function $f = x_1^{-c} ((a - c) x_3 - b x_1)^a$ as solution. This pattern occurs for all the algebras in this case, up to $\mathfrak{g}_{6,79}$.

3.7 Nilradical isomorphic to $\mathfrak{g}_{5,4}$

There are 12 isomorphism classes having this algebra as nilradical, eight of them depending on one or more parameters [tables 13 and 14]. This is also the first case where the number of invariants depends essentially on the action of the vector X_6 on the nilradical $\mathfrak{g}_{5,4}$. As follows from [8], the structure of solvable Lie algebras having this algebra as nilradical is

$$\begin{aligned} [X_2, X_4] &= X_1; [X_3, X_5] = X_1 \\ [X_i, X_6] &= C_{i6}^k x_k, \quad 1 \leq i, k \leq 5 \end{aligned}$$

Therefore the matrix $(C_{ij}^k x_k)$ representing the commutator table is

$$A = \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & C_{16}^k x_k \\ 0 & 0 & 0 & x_1 & 0 & C_{26}^k x_k \\ 0 & 0 & 0 & 0 & x_1 & C_{36}^k x_k \\ 0 & -x_1 & 0 & 0 & 0 & C_{46}^k x_k \\ 0 & 0 & -x_1 & 0 & 0 & C_{56}^k x_k \\ -C_{16}^k x_k & -C_{26}^k x_k & -C_{36}^k x_k & -C_{46}^k x_k & -C_{56}^k x_k & 0 \end{pmatrix} \quad (12)$$

whose determinant is $d = x_1^4 (C_{16}^k x_k)^2$. Therefore, we will obtain non-trivial invariants if and only if $C_{16}^k = 0$ for $1 \leq k \leq 5$. If this occurs, A is easily seen to have rank four and by the Beltrametti-Blasi formula [19] the algebra will have exactly two invariants. Moreover, since this implies that X_1 is a central element, one invariant can be chosen as $I_1 = x_1$.

Lemma 1 *Let $\mathfrak{g}$ be a solvable Lie algebra whose nilradical is isomorphic to $\mathfrak{g}_{5,4}$. If $\det(A) = 0$, then $\mathfrak{g}$ admits a*

fundamental set of invariants formed by two polynomial functions, $I_1 = x_1$ being one of them.

Thus in this case we obtain classical Casimir operators [20, 23]. Observe further that $\mathfrak{g}_{5,4}$ is isomorphic to the five dimensional Heisenberg Lie algebra. In [25] the invariants of solvable algebras having a Heisenberg Lie algebra as nilradical are determined generically.

3.8 Nilradical isomorphic to $\mathfrak{g}_{5,5}$

Five isomorphism classes arise from Mubarakzhanovs classification [8] for this nilradical, only one of them, $\mathfrak{g}_{6,94}$, depending on a parameter [table 15]. An argumentation like in the preceding case shows that these algebras have no non-trivial invariants, up to the algebra $\mathfrak{g}_{6,94}$ for the value $\lambda + 2 = 0$, for which we obtain again a fundamental set of invariants formed by two polynomial functions.

3.9 Nilradical isomorphic to $\mathfrak{g}_{5,6}$

Finally, for the nilradical $\mathfrak{g}_{5,6}$ there is a unique non-parametrized isomorphism class, $\mathfrak{g}_{6,99}$ [table 16]. For this algebra is easily verified that the matrix $(C_{ij}^k x_k)$ has maximal rank, which shows that this algebra has no invariants.

The last three nilradicals, $\mathfrak{g}_{5,4}$, $\mathfrak{g}_{5,5}$ and $\mathfrak{g}_{5,6}$ have in common that they are all deformations of the 5-dimensional Heisenberg Lie algebra (considering the trivial deformation for the Lie algebra $\mathfrak{g}_{5,4}$)

[36]. This fact allows us to deduce a similar result which holds in arbitrary dimension: For $m \leq 2$ let $\mathfrak{n}_{\alpha_1, \dots, \alpha_{\lfloor \frac{m}{2} \rfloor + 2}}$ be the $(2m+1)$ -dimensional nilpotent Lie algebra given by the brackets

$$\left. \begin{aligned} [X_{2+j}, X_{2m+1-j}] &= X_1, & 0 \leq j \leq m-1 \\ [X_{2+j}, X_{2m+1}] &= \alpha_j X_{1+j}, & 1 \leq j \leq 2m-2 \\ \alpha_j + \alpha_{2m-j} &= 0, & 1 \leq j \leq m-1 \end{aligned} \right\}, \quad (13)$$

over the basis $\{X_1, \dots, X_{2m+1}\}$. Let $\mathfrak{r}_m$ be a $(2m+2)$ -dimensional solvable Lie algebra whose nilradical is isomorphic to $\mathfrak{n}_{\alpha_1, \dots, \alpha_{\lfloor \frac{m}{2} \rfloor + 2}}$.

Theorem 3 *Let $\mathfrak{r}_m$ be a $(2m+2)$ -dimensional solvable Lie algebra whose nilradical is isomorphic to $\mathfrak{n}_{\alpha_1, \dots, \alpha_{\lfloor \frac{m}{2} \rfloor + 2}}$.*

1. *If the action $\left(\frac{r_m}{\mathfrak{n}_{\alpha_1, \dots, \alpha_{\lfloor \frac{m}{2} \rfloor + 2}}} \right)$ over X_1 is nonzero, then $\mathcal{N}(\mathfrak{r}_m) = 0$.*
2. *If $\left(\frac{r_m}{\mathfrak{n}_{\alpha_1, \dots, \alpha_{\lfloor \frac{m}{2} \rfloor + 2}}} \right) \cdot X_1 = 0$, then $\mathfrak{r}_m$ admits a fundamental set of invariants formed by two elements, one of which can be taken as $I_1 = x_1$.*

Proof. If $(C_{ij}^k x_k)$ is the matrix associated to the commutators of $\mathfrak{r}_m$, then it can easily be proven by induction over m that

$$\det(C_{ij}^k) = x_1^{2m} (C_{1, 2m+2}^k x_k)^2$$

so that $\mathcal{N}(\mathfrak{r}_m) = 0$ if and only if $C_{1, 2m+2}^k = 0$ for all k . Otherwise we obtain that $\text{rank}(C_{ij}^k) = 2m$, thus the algebra has two invariants. The second assertion follows from the fact that X_1 spans the center of the nilradical. ■

Although there is no procedure to determine the detailed structure of the second invariant, the analysis of low values of m seems to indicate that it can always be chosen as a polynomial function. The exact structure of this polynomial depends heavily on the parameters α_i , and cannot therefore be comprised in a formula.

Acknowledgement 1 *The authors wishes to express his gratitude to P. Turkowski for providing him with copies of the articles [7] and [33], as well as to the Ramon Areces Foundation for support during this research.*

References

- [1] H. Casimir. Über die Konstruktion einer zu den irreduziblen Darstellungen halbeinfacher kontinuierlicher Gruppen gehörigen Differentialgleichung, Proc. Roy. Acad. Amsterdam **34** (1931), 844-846.
- [2] Ch. Yen. Sur les polynomes de Poincaré des groupes simples exceptionnels, C. R. Acad. Sci. Paris **228** (1949), 628-630.

- [3] W. Miller. Lie theory and special functions, Academic, New York 1968.
- [4] M. Gell-Mann. Symmetries of baryons and mesons, Phys. Rev. **125** (1962), 1067-1084.
- [5] J. Patera, R. T. Sharp, P. Winternitz and H. Zassenhaus. Invariants of real low dimension Lie algebras, J. Mathematical Phys. **17** (1976), 986-994.
- [6] J. C. Ndogmo. Invariants of solvable Lie algebras of dimension six. J. Phys. A **33** (2000), 2273-2287.
- [7] P. Turkowski. Solvable Lie algebras of dimension six. J. Math. Phys. **31** (1990), 1344-1350.
- [8] G. M. Mubarakzhanov. Classification of solvable Lie algebras of sixth order with a non-nilpotent basis element (Russian), Izv. Vysš. Učebn. Zaved. Matematika **35** (1963), 104-116.
- [9] A. A. Arhangelsky. Completely integrable hamiltonian systems on a group of triangular matrices, Mat. Sb. **108** (1979), 134-142.
- [10] V. V. Trofimov. Semi-invariants of a coadjoint representation of Borel subalgebras of simple Lie algebras, Trudy Sem. Vektor Tenzor. Anal. **21** (1983), 84-105.
- [11] M. Pauri, G. M. Prosperi. On the construction of the invariant operators for any finite-parameter Lie group, Nuovo Cimento, X. Ser. A **43** (1966), 533-537.
- [12] R. Mirman. Number of polynomial invariants of adjoint and fundamental compact inhomogeneous unitary algebras, J. Math. Phys. **9** (1968), 47-49.
- [13] A. T. Fomenko, V. V. Trofimov. Integrable systems on Lie algebras and symmetric spaces, Gordon and Breach, New York 1989.
- [14] J. M. Levy-Leblond. Group theory and its applications, New York: Academic Press 1972.
- [15] P. Roman, J. J. Aghassi and P. L. Huddleston. Relativistic dynamical symmetries and dilatations, J. Mathematical Phys. **13** (1972), 1852-1861.
- [16] V. V. Trofimov. Euler equations on Borel subalgebras of semisimple Lie algebras, Izv. Akad. Nauk. SSSR Ser. Mat. **43** (1979), 714-732.
- [17] G. Racah. Sulla caratterizzazione delle rappresentazioni irriducibili dei gruppi semisemplici di Lie. (Italian) Atti Accad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Nat. **8** (1950), 108-112.
- [18] Ch. Schweigert and D. Fuchs. Symmetries, Lie algebras and representations, Cambridge Univ. Press 1997.
- [19] E. G. Beltrametti and A. Blasi. On the number of Casimir operators associated with any Lie group, Phys. Lett. **20** (1966), 62-64.
- [20] L. Abellanas and L. Martínez Alonso. A general setting for Casimir invariants, J. Mathematical Phys. **16** (1975), 1580-1584.

- [21] L. Solomon, D.-N. Verma. Sur le corps des quotients de l'algèbre enveloppante d'une algèbre de Lie, C. R. Acad. Sci. Paris Sé9r. A-B **264** (1967) 985-986.
- [22] J. E. Humphreys. Linear algebraic groups. Springer-Verlag, New York-Heidelberg, 1975.
- [23] J. Dixmier. Algèbres enveloppantes. Hermann, Paris 1974.
- [24] J. L. Rubin, P. Winternitz. Solvable Lie algebras with Heisenberg ideals. J. Phys. A **26** (1993), 1123-1138.
- [25] J. C. Ndogmo, P. Winternitz. Generalized Casimir operators of solvable Lie algebras with abelian nilradicals. J. Phys. A **27** (1994), 2787-2800.
- [26] R. Campoamor-Stursberg. Invariants of solvable rigid Lie algebras up to dimension 8, J. Phys. A **35** (2002), 6293-6306.
- [27] R. Campoamor-Stursberg. On the invariants of some solvable rigid Lie algebras, J. Math. Phys., to appear.
- [28] E. L. Dixon. Differential equations from the group standpoint, Annals of Math. **25** (1924), 287-378.
- [29] G. M. Mubarakzyanov. Some theorems on solvable Lie algebras, Sb. Asp. Rabot. Mat. Mech. Phys. (1963), 103.
- [30] G. M. Mubarakzyanov. Classification of real structures of Lie algebras of fifth order (Russian), Izv. Vysš. Učebn. Zaved. Matematika **34** (1963), 99-106.
- [31] G. M. Mubarakzyanov. On solvable Lie algebras (Russian), Izv. Vysš. Učehn. Zaved. Matematika **32** (1963), 114-123.
- [32] V. V. Morozov. Classification of nilpotent Lie algebras of sixth order (Russian), Izv. Vysš. Učebn. Zaved. Matematika **5** (1958), 161-171.
- [33] P. Turkowski. Literature on the structure of low dimensional nonsemisimple Lie algebras and its applications to cosmology, Acta Cosmologica **20** (1994), 147-153.
- [34] J. C. Ndogmo. Properties of the invariants of solvable Lie algebras, Canad. Math. Bull. **43** (2000), 459-471.
- [35] P. Olver. Applications of Lie groups to differential equations, Springer-Verlag, New York 1986.
- [36] R. Campoamor-Stursberg. On k -abelian p -filiform Lie algebras I, Acta Math. Univ. Comen. **71** (2002), 51-68.

Table 2: Nilradical isomorphic to $5\mathfrak{g}_1$

Name	Brackets		Invariants
$\mathfrak{g}_{6,1}$ $0 < \delta \leq \gamma $ $\leq \beta \leq \alpha < 1$	$[X_1, X_6] = X_1$ $[X_3, X_6] = \beta X_3$ $[X_5, X_6] = \delta X_5$	$[X_2, X_6] = \alpha X_2$ $[X_4, X_6] = \gamma X_4$	$I_1 = x_2 x_1^{-\alpha}$ $I_2 = x_3 x_1^{-\beta}$ $I_3 = x_4 x_1^{-\gamma}$ $I_4 = x_5 x_1^{-\delta}$
$\mathfrak{g}_{6,2}$ $0 < \delta \leq \gamma \leq 1$	$[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = X_3$ $[X_5, X_6] = \delta X_5$	$[X_2, X_6] = X_1 + \alpha X_2$ $[X_4, X_6] = \gamma X_4$	$I_1 = x_1 \exp\left(\frac{-\alpha x_2}{x_1}\right)$ $I_2 = x_3 \exp\left(\frac{-x_2}{x_1}\right)$ $I_3 = x_4 \exp\left(\frac{-\gamma x_2}{x_1}\right)$ $I_4 = x_5 \exp\left(\frac{-\delta x_2}{x_1}\right)$
$\mathfrak{g}_{6,3}$ $0 < \delta \leq 1$	$[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = X_2 + \alpha X_3$ $[X_5, X_6] = \delta X_5$	$[X_2, X_6] = X_1 + \alpha X_2$ $[X_4, X_6] = X_4$	$I_1 = x_1 \exp\left(\frac{-\alpha x_2}{x_1}\right)$ $I_2 = \frac{2x_1 x_3 - x_2^2}{x_1^2}$ $I_3 = x_4 \exp\left(\frac{-x_2}{x_1}\right)$ $I_4 = x_5 \exp\left(\frac{-\delta x_2}{x_1}\right)$
$\mathfrak{g}_{6,4}$	$[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = X_2 + \alpha X_3$ $[X_5, X_6] = X_5$	$[X_2, X_6] = X_1 + \alpha X_2$ $[X_4, X_6] = X_3 + \alpha X_4$	$I_1 = x_1 \exp\left(\frac{-\alpha x_2}{x_1}\right)$ $I_2 = \frac{(2x_1 x_3 - x_2^2)}{x_1^2}$ $I_3 = \frac{(3x_1^2 x_4 + x_2^3 - 3x_1 x_2 x_3)}{x_1^2}$ $I_4 = x_5 \exp\left(\frac{-x_2}{x_1}\right)$
$\mathfrak{g}_{6,5}$	$[X_1, X_6] = X_1$ $[X_3, X_6] = X_2 + X_3$ $[X_5, X_6] = X_4 + X_5$	$[X_2, X_6] = X_1 + X_2$ $[X_4, X_6] = X_3 + X_4$	$I_1 = x_1 \exp\left(\frac{-x_2}{x_1}\right)$ $I_2 = \frac{(2x_1 x_3 - x_2^2)}{x_1^2}$ $I_3 = \frac{(3x_1^2 x_4 + x_2^3 - 3x_1 x_2 x_3)}{x_1^2}$ $I_4 = \frac{x_2^4 - 8x_1^3 x_5 - 4x_1 x_2^2 x_3 + 8x_1^2 x_2 x_4}{x_1^4}$
$\mathfrak{g}_{6,6}$ $\alpha \leq \beta$	$[X_1, X_6] = X_1$ $[X_3, X_6] = X_2 + \alpha X_3$ $[X_5, X_6] = X_4 + \beta X_5$	$[X_2, X_6] = \alpha X_2$ $[X_4, X_6] = \beta X_4$	$I_1 = x_2 x_1^{-\alpha}$ $I_2 = x_1 \exp\left(\frac{-x_3}{x_2}\right)$ $I_3 = x_4 x_1^{-\beta}$ $I_4 = x_1 \exp\left(\frac{-x_5}{x_4}\right)$
$\mathfrak{g}_{6,7}$ $\alpha^2 + \beta^2 \neq 0$	$[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = X_2 + \alpha X_3$ $[X_5, X_6] = X_4 + \beta X_5$	$[X_2, X_6] = X_1 + \alpha X_2$ $[X_4, X_6] = \beta X_4$	$I_1 = x_1 \exp\left(\frac{-\alpha x_2}{x_1}\right)$ $I_2 = \frac{(2x_1 x_3 - x_2^2)}{x_1^2}$ $I_3 = \frac{(x_1 x_5 - x_2 x_4)}{x_1} \exp\left(\frac{-\beta x_2}{x_1}\right)$ $I_4 = x_4 \exp\left(\frac{-\beta x_2}{x_1}\right)$
$\mathfrak{g}_{6,8}$ $0 < \gamma \leq \beta \leq \alpha $	$[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = \gamma X_3$ $[X_5, X_6] = X_4 + p X_5$	$[X_2, X_6] = \beta X_2$ $[X_4, X_6] = p X_4 - X_5$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 + i x_5}{x_4 - i x_5}\right)^{ip}$ $I_2 = x_1 \exp^{\alpha \arctan x_4 x_5^{-1}}$ $I_3 = x_2 \exp^{\beta \arctan x_4 x_5^{-1}}$ $I_4 = x_3 \exp^{\gamma \arctan x_4 x_5^{-1}}$

Table 3: Nilradical isomorphic to $5\mathfrak{g}_1$ (cont.)

Name	Brackets		Invariants
$\mathfrak{g}_{6,9}$	$[X_1, X_6] = \alpha X_1$	$[X_2, X_6] = \beta X_2$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 + ix_5}{x_4 - ix_5} \right)^{ip}$
$\alpha \neq 0$	$[X_3, X_6] = X_2 + \beta X_3$	$[X_4, X_6] = pX_4 - X_5$	$I_2 = x_1 \exp^{\alpha \arctan x_4 x_5^{-1}}$
	$[X_5, X_6] = X_4 + pX_5$		$I_3 = x_2 \exp^{\beta \arctan x_4 x_5^{-1}}$
			$I_4 = (x_3 + x_2 \arctan \frac{x_4}{x_5}) \exp^{\beta \arctan \frac{x_4}{x_5}}$
$\mathfrak{g}_{6,10}$	$[X_1, X_6] = \alpha X_1$	$[X_2, X_6] = X_1 + \alpha X_2$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 + ix_5}{x_4 - ix_5} \right)^{ip}$
	$[X_3, X_6] = X_2 + \alpha X_3$	$[X_4, X_6] = pX_4 - X_5$	$I_2 = x_1 \exp^{\alpha \arctan x_4 x_5^{-1}}$
	$[X_5, X_6] = X_4 + pX_5$		$I_3 = (x_2 + x_1 \arctan \frac{x_4}{x_5}) \exp^{\alpha \arctan \frac{x_4}{x_5}}$
			$I_4 = \frac{2x_3 x_1 - x_2^2}{x_1}$
$\mathfrak{g}_{6,11}$	$[X_1, X_6] = \alpha X_1$	$[X_2, X_6] = pX_2 - X_3$	$I_1 = (x_2^2 + x_3^2) \left(\frac{x_2 + ix_3}{x_2 - ix_3} \right)^{ip}$
$\alpha s \neq 0$	$[X_3, X_6] = X_2 + pX_3$	$[X_4, X_6] = qX_4 - sX_5$	$I_2 = (x_4^2 + x_5^2) \left(\frac{x_4 + ix_5}{x_4 - ix_5} \right)^{iqs^{-1}}$
	$[X_5, X_6] = sX_4 + qX_5$		$I_3 = (x_2^2 + x_3^2)^\alpha x_1^{-2p}$
			$I_4 = (x_4^2 + x_5^2)^\alpha x_1^{-2q}$
$\mathfrak{g}_{6,12}$	$[X_1, X_6] = \alpha X_1$	$[X_2, X_6] = pX_2 - X_3$	$I_1 = (x_3 + ix_2)^\alpha x_1^{(i-p)}$
$\alpha \neq 0$	$[X_3, X_6] = X_2 + pX_3$	$[X_4, X_6] = X_2 + pX_4 - X_5$	$I_2 = \frac{((ix_4 + x_5)\alpha - (x_3 + ix_2) \ln(x_1))^\alpha}{x_1^{p-i}}$
	$[X_5, X_6] = X_3 + X_4 + pX_5$		$I_3 = (x_3 - ix_2)^\alpha x_1^{p+i}$
			$I_4 = \frac{((-ix_4 + x_5) - (x_3 - ix_2) \ln(x_1))^\alpha}{x_1^{p+i}}$

Table 4: Nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$

Name	Brackets		Invariants
$\mathfrak{g}_{6,13}$ $\alpha \neq 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (\alpha + \beta) X_1$	$I_1 = x_4^{(\alpha+\beta)} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = \beta X_3$	$I_2 = x_4^h x_5^{-1}$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = h X_5$	
$\mathfrak{g}_{6,14}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (\alpha + \beta) X_1$	$I_1 = x_4^{(\alpha+\beta)} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = \beta X_3$	$I_2 = x_1 \exp\left(\frac{-(\alpha+\beta)x_5}{x_1}\right)$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = X_1 + (\alpha + \beta) X_5$	
$\mathfrak{g}_{6,15}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (1 + h) X_1$	$I_1 = x_4^{(1+h)} x_1^{-1}$
	$[X_2, X_6] = X_2 + X_4$	$[X_3, X_6] = h X_3 + X_5$	$I_2 = x_5 x_4^{-h}$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = h X_5$	
$\mathfrak{g}_{6,16}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = X_1$	$I_1 = x_5$
	$[X_2, X_6] = X_2 + X_4$	$[X_3, X_6] = X_5$	$I_2 = x_1 \exp\left(\frac{-x_4}{x_1}\right)$
	$[X_4, X_6] = X_1 + X_4$		
$\mathfrak{g}_{6,17}$ $\alpha^2 + \varepsilon^2 \neq 0$ $\alpha\varepsilon = 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = \alpha X_1$	$I_1 = x_5^\alpha x_1^{-1}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = X_4$	$I_2 = \varepsilon x_1 - \alpha x_4$
	$[X_4, X_6] = \varepsilon X_1$	$[X_5, X_6] = X_5$	$(\alpha = 0) : = x_5 \exp^{-x_4(\varepsilon x_1)^{-1}}$
$\mathfrak{g}_{6,18}$ $\beta \neq 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (1 + \alpha) X_1$	$I_1 = x_4^{(1+\alpha)} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = X_3 + X_4$	$I_2 = x_5 x_4^{-\beta}$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = \beta X_5$	
$\mathfrak{g}_{6,19}$ $\beta \neq 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (1 + \alpha) X_1$	$I_1 = x_4^{(1+\alpha)} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = X_3 + X_4$	$I_2 = x_4 \exp\left(\frac{-x_5}{x_1}\right)$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = X_1 + (\alpha + 1) X_5$	
$\mathfrak{g}_{6,20}$ $\beta \neq 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = X_1$	$I_1 = x_5 x_1^{-\beta}$
	$[X_2, X_6] = 0$	$[X_3, X_6] = X_3 + X_4$	$I_2 = x_1 \exp\left(\frac{-x_4}{x_1}\right)$
	$[X_4, X_6] = X_1 + X_4$	$[X_5, X_6] = \beta X_5$	
$\mathfrak{g}_{6,21}$ $h \neq 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2\alpha X_1$	$I_1 = x_4^{2\alpha} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2 + X_3$	$[X_3, X_6] = \alpha X_3$	$I_2 = x_5 x_4^{-h}$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = h X_5$	
$\mathfrak{g}_{6,22}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2\alpha X_1$	$I_1 = x_4^{2\alpha} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2 + X_3$	$[X_3, X_6] = \alpha X_3$	$I_2 = x_4 \exp\left(\frac{-x_5}{x_1}\right)$
	$[X_4, X_6] = X_4$	$[X_5, X_6] = X_1 + 2\alpha X_5$	
$\mathfrak{g}_{6,23}$ $\varepsilon h = 0$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2\alpha X_1$	$I_1 = x_4^{2\alpha} x_1^{-1}$
	$[X_2, X_6] = \alpha X_2 + X_3$	$[X_3, X_6] = \alpha X_3 + X_4$	$I_2 = \frac{(h x_5 + \varepsilon x_1)^{2\alpha}}{x_1^{2\alpha+h}} \quad (h \neq 0)$
	$[X_4, X_6] = \alpha X_4$	$[X_5, X_6] = \varepsilon X_1 + (2\alpha + h) X_5$	$= \frac{2\alpha x_5 - \varepsilon x_1 \ln(x_1)}{x_1} \quad (h = 0)$

Table 5: Nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$ (cont.)

Name	Brackets	Invariants
$\mathfrak{g}_{6,24}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 0$ $I_1 = x_1$
	$[X_2, X_6] = X_3$	$[X_3, X_6] = X_4$ $I_2 = x_5^h \exp\left(\frac{-x_4}{x_1}\right)$
	$[X_4, X_6] = hX_1$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,25}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (1+h)X_1$ $I_1 = x_1^{-\beta} x_5^{1+h}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = hX_3$ $I_2 = x_1 \exp\left(\frac{-(1+h)x_4}{x_5}\right)$
	$[X_4, X_6] = \beta X_4 + X_5$	$[X_5, X_6] = \beta X_5$
$\mathfrak{g}_{6,26}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (1+h)X_1$ $I_1 = \frac{2x_1x_4 - x_5^2}{x_1^2}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = hX_3$ $I_2 = x_1 \exp\left(\frac{-(1+h)x_5}{x_1}\right)$
	$[X_4, X_6] = (1+h)X_4 + X_5$	$[X_5, X_6] = X_1 + (1+h)X_5$
$\mathfrak{g}_{6,27}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (\alpha + \beta)X_1$ $I_1 = \frac{(\alpha x_5 - \varepsilon x_1)^{\alpha+\beta}}{x_1^{\alpha+\beta}} x_1^\beta$
$\alpha^2 + \beta^2 \neq 0$	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = \beta X_3 + X_4$ $I_2 = x_1^{\frac{\alpha+\beta}{\alpha}} \exp\left(\frac{\alpha^2 x_4 - \varepsilon x_1}{\alpha x_5 - \varepsilon x_1}\right)$
$\alpha \varepsilon = 0$	$[X_4, X_6] = \beta X_4 + X_5$	$[X_5, X_6] = \varepsilon X_4 + \beta X_5$
$\mathfrak{g}_{6,27}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 0$ $I_1 = x_1$
$\alpha + \beta = 0$	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = -\alpha X_3 + X_4$ $I_2 = \frac{\alpha x_5 - \varepsilon x_1}{x_1} \exp\left(\frac{\alpha^2 x_4 - \varepsilon x_1}{\alpha x_5 - \varepsilon x_1}\right)$
$\alpha \varepsilon = 0$	$[X_4, X_6] = -\alpha X_4 + X_5$	$[X_5, X_6] = \varepsilon X_4 - \alpha X_5$
$\mathfrak{g}_{6,28}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2X_1$ $I_1 = x_5^2 x_1^{-\beta}$
	$[X_2, X_6] = X_2 + X_3$	$[X_3, X_6] = X_3$ $I_2 = x_1 \exp\left(\frac{-\alpha x_4}{x_5}\right)$
	$[X_4, X_6] = \beta X_4 + X_5$	$[X_5, X_6] = \beta X_5$
$\mathfrak{g}_{6,29}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = (\alpha + \beta)X_1$ $I_1 = \frac{-2x_1x_4 + x_5^2}{x_1^2}$
	$[X_2, X_6] = \alpha X_2$	$[X_3, X_6] = \beta X_3 + X_4$ $I_2 = x_1 \exp\left(\frac{-2x_5}{x_1}\right)$
	$[X_4, X_6] = \beta X_4 + X_5$	$[X_5, X_6] = \varepsilon X_4 + \beta X_5$
$\mathfrak{g}_{6,30}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 0$ $I_1 = x_1$
	$[X_2, X_6] = X_3$	$[X_3, X_6] = 0$ $I_2 = x_5 \exp\left(\frac{-x_4}{x_5}\right)$
	$[X_4, X_6] = X_4 + X_5$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,31}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2X_1$ $I_1 = \frac{-2x_1x_4 + x_5^2}{x_1^2}$
	$[X_2, X_6] = X_2 + X_3$	$[X_3, X_6] = X_3 + X_4$ $I_2 = x_1 \exp\left(\frac{-2x_4}{x_5}\right)$
	$[X_4, X_6] = X_4 + X_5$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,32}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2\alpha X_1$ $I_1 = x_5^{2\alpha} x_1^{-\gamma}$
$2\alpha + h > \gamma$	$[X_2, X_6] = \alpha X_2 + X_3$	$[X_3, X_6] = -X_2 + \alpha X_3$ $I_2 = (hx_4 + \varepsilon x_1)^\gamma x_5^{-(2\alpha+h)}$
$\varepsilon h = 0$	$[X_4, X_6] = \varepsilon X_1 + (2\alpha + h)X_4$	$[X_5, X_6] = \gamma X_5$
$\mathfrak{g}_{6,33}$	$[X_2, X_3] = X_1$	$[X_1, X_6] = 2\alpha X_1$ $I_1 = x_4^{2\alpha} x_1^{-\beta}$
$\beta \leq 2\alpha$	$[X_2, X_6] = \alpha X_2 + X_3$	$[X_3, X_6] = -X_2 + \alpha X_3$ $I_2 = x_1 \exp\left(\frac{-2\alpha x_5}{x_1}\right)$
	$[X_4, X_6] = \beta X_4$	$[X_5, X_6] = X_1 + 2\alpha X_5$

Table 6: Nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$ (cont.)

Name	Brackets	Invariants
$\mathfrak{g}_{6,34}$	$[X_2, X_3] = X_1$	$I_1 =$
$\varepsilon h = 0$	$[X_2, X_6] = \alpha X_2 + X_3$	$I_2 =$
	$[X_4, X_6] = (2\alpha + h) X_4 + X_5$	$[X_1, X_6] = 2\alpha X_1$
		$[X_3, X_6] = -X_2 + \alpha X_3$
		$[X_5, X_6] = (2\alpha + h) X_5 + \varepsilon X_1$
$\mathfrak{g}_{6,35}$	$[X_2, X_3] = X_1$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 - ix_5}{x_4 + ix_5} \right)^{i\gamma}$
$\alpha^2 + \beta^2 \neq 0$	$[X_2, X_6] = \alpha X_2$	$I_2 = x_1 \exp \left(-(\alpha + \beta) \arctan \left(\frac{x_4}{x_5} \right) \right)$
	$[X_4, X_6] = \gamma X_4 + X_5$	$[X_3, X_6] = \beta X_3$
		$[X_5, X_6] = -X_4 + \gamma X_5$
$\mathfrak{g}_{6,36}$	$[X_2, X_3] = X_1$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 - ix_5}{x_4 + ix_5} \right)^{i\gamma}$
	$[X_2, X_6] = \alpha X_2 + X_3$	$I_2 = x_1 \exp \left(-2\alpha \arctan \left(\frac{x_4}{x_5} \right) \right)$
	$[X_4, X_6] = \gamma X_4 + X_5$	$[X_3, X_6] = \alpha X_3$
		$[X_5, X_6] = -X_4 + \gamma X_5$
$\mathfrak{g}_{6,37}$	$[X_2, X_3] = X_1$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 - ix_5}{x_4 + ix_5} \right)^{\frac{\beta i}{s}}$
$s \neq 0$	$[X_2, X_6] = \alpha X_2 + X_3$	$I_2 = x_1 \exp \left(-\frac{s}{2} \alpha \arctan \left(\frac{x_4}{x_5} \right) \right)$
	$[X_4, X_6] = \beta X_4 + s X_5$	$[X_3, X_6] = -X_2 + \alpha X_3$
		$[X_5, X_6] = -s X_4 + \beta X_5$
$\mathfrak{g}_{6,38}$	$[X_2, X_3] = X_1$	$I_1 = (x_4^2 + x_5^2) \left(\frac{x_4 - ix_5}{x_4 + ix_5} \right)^{i\alpha}$
	$[X_2, X_6] = \alpha X_2 + X_3 + X_4$	$I_2 = x_1 \exp \left(-2\alpha \arctan \left(\frac{x_4}{x_5} \right) \right)$
	$[X_4, X_6] = \alpha X_4 + X_5$	$[X_3, X_6] = -X_2 + \alpha X_3 + X_5$
		$[X_5, X_6] = -X_4 + \alpha X_5$

Table 7: Nilradical isomorphic to $\mathfrak{g}_{4,1} + \mathfrak{g}_1$

Name	Brackets		Invariants
$\mathfrak{g}_{6,39}$ $\gamma \neq 0$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3^{2+h} x_2^{-\gamma}$
	$[X_1, X_6] = (1+h) X_1$	$[X_2, X_6] = (2+h) X_2$	$I_2 = \frac{x_1^2 - 2x_2 x_4}{x_1^2 - 2x_2 x_4} x_2 x_3 \frac{h}{\gamma}$
	$[X_3, X_6] = \gamma X_3$	$[X_4, X_6] = h X_4$	
	$[X_5, X_6] = X_5$		
$\mathfrak{g}_{6,40}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_2 \exp\left(\frac{-(2+h)x_3}{x_2}\right)$
	$[X_1, X_6] = (1+h) X_1$	$[X_2, X_6] = (2+h) X_2$	$I_2 = \frac{x_1^2 - 2x_2 x_4}{x_2} \exp\left(\frac{-h x_3}{x_2}\right)$
	$[X_3, X_6] = X_2 + (2+h) X_3$	$[X_4, X_6] = h X_4$	
	$[X_5, X_6] = X_5$		
$\mathfrak{g}_{6,41}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3^{2+h} x_2^{-h}$
	$[X_1, X_6] = (1+h) X_1$	$[X_2, X_6] = (2+h) X_2$	$I_2 = (x_2 x_3)^{-1} \exp\left(\frac{(1+h)(2x_2 x_4 - x_1^2)}{x_2 x_3}\right)$
	$[X_3, X_6] = h X_3$	$[X_4, X_6] = X_3 + h X_4$	
	$[X_5, X_6] = X_5$		
$\mathfrak{g}_{6,42}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3^{2+h} x_2^{-1}$
	$[X_1, X_6] = (1+h) X_1$	$[X_2, X_6] = (2+h) X_2$	$I_2 = \frac{(2x_2 x_4 - x_1^2)}{x_2 x_3^h}$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = h X_4$	
	$[X_5, X_6] = X_3 + X_5$		
$\mathfrak{g}_{6,43}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_1^2 - 2x_2 x_4$
	$[X_1, X_6] = 0$	$[X_2, X_6] = X_2$	$I_2 = x_2 \exp\left(\frac{-x_3}{x_2}\right)$
	$[X_3, X_6] = X_2 + X_3$	$[X_4, X_6] = -X_4$	
	$[X_5, X_6] = X_3 + X_5$		
$\mathfrak{g}_{6,44}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3^3 x_2^{-\gamma}$
	$[X_1, X_6] = 2X_1$	$[X_2, X_6] = 3X_2$	$I_2 = \frac{(2x_2 x_4 - x_1^2)^3}{x_2^4}$
	$[X_3, X_6] = \gamma X_3$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = X_4 + X_5$		
$\mathfrak{g}_{6,45}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_2 \exp\left(\frac{-3x_3}{x_2}\right)$
	$[X_1, X_6] = 2X_1$	$[X_2, X_6] = 3X_2$	$I_2 = \frac{(2x_2 x_4 - x_1^2)^3}{x_2^4}$
	$[X_3, X_6] = X_2 + 3X_3$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = X_4 + X_5$		
$\mathfrak{g}_{6,46}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3 \exp\left(\frac{x_1^2 - 2x_2 x_4}{2x_2 x_3}\right)$
	$[X_1, X_6] = 2X_1$	$[X_2, X_6] = 3X_2$	$I_2 = \frac{x_3^3}{x_2}$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = X_3 + X_4$	
	$[X_5, X_6] = X_4 + X_5$		
$\mathfrak{g}_{6,47}$ $\gamma \neq 0$ $\epsilon \neq 0, \pm 1$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_3 x_2^{-\gamma}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = X_2$	$I_2 = x_2^\epsilon \exp\left(\frac{x_1^2 - 2x_2 x_4}{x_2} 2x_2^2\right)$
	$[X_3, X_6] = \gamma X_3$	$[X_4, X_6] = \epsilon X_2 + X_4$	
	$[X_5, X_6] = 0$		
$\mathfrak{g}_{6,48}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$	$I_1 = x_2 \exp\left(\frac{-x_3}{x_2}\right)$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = X_2$	$I_2 = \frac{2x_2 x_4 - x_1^2}{x_2^2}$
	$[X_3, X_6] = X_2 + X_3$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = 0$		

Table 8: Nilradical isomorphic to $\mathfrak{g}_{3,1} + 2\mathfrak{g}_1$ (cont.)

Name	Brackets	Invariants
$\mathfrak{g}_{6,49}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$
$\epsilon \neq 0, \pm 1$	$[X_1, X_6] = X_1$	$[X_2, X_6] = X_2$
	$[X_3, X_6] = 0$	$[X_4, X_6] = \epsilon X_2 + X_4$
	$[X_5, X_6] = X_3$	
$\mathfrak{g}_{6,50}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$
$\epsilon \neq 0, \pm 1$	$[X_1, X_6] = X_1$	$[X_2, X_6] = X_2$
	$[X_3, X_6] = \epsilon X_2 + X_3$	$[X_4, X_6] = X_3 + X_4$
	$[X_5, X_6] = 0$	
$\mathfrak{g}_{6,51}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$
$\epsilon \neq \pm 1$	$[X_1, X_6] = 0$	$[X_2, X_6] = 0$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = \epsilon X_2$
	$[X_5, X_6] = 0$	
$\mathfrak{g}_{6,52}$	$[X_1, X_5] = X_2$	$[X_4, X_5] = X_1$
$\epsilon \neq 0, \pm 1$	$[X_1, X_6] = 0$	$[X_2, X_6] = 0$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = \epsilon X_2$
	$[X_5, X_6] = X_4$	

Table 9: Nilradical isomorphic to $\mathfrak{g}_{5,1}$

Name	Brackets		Invariants
$\mathfrak{g}_{6,53}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_1$
	$[X_1, X_6] = 0$	$[X_2, X_6] = 0$	$I_2 = x_2$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = -X_5$		
$\mathfrak{g}_{6,54}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-\lambda}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = \lambda X_2$	$I_2 = (x_1 x_4 - x_2 x_3) x_1^{\alpha+1-\gamma}$
	$[X_3, X_6] = (1-\gamma) X_3$	$[X_4, X_6] = (\lambda-\gamma) X_4$	
	$[X_5, X_6] = \gamma X_5$		
$\mathfrak{g}_{6,55}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-(1+\gamma)}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = (1+\gamma) X_2$	$I_2 = x_1 \exp\left(\frac{x_2 x_3 - x_1 x_4}{x_1^2}\right)$
	$[X_3, X_6] = (1-\gamma) X_3$	$[X_4, X_6] = X_1 + X_4$	
	$[X_5, X_6] = \gamma X_5$		
$\mathfrak{g}_{6,56}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-(1-\gamma)}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = (1-\gamma) X_2$	$I_2 = x_1^{\gamma-1} \exp\left(\frac{(1-\gamma)(x_2 x_3 - x_1 x_4)}{x_2^2}\right)$
	$[X_3, X_6] = X_2 + (1-\gamma) X_3$	$[X_4, X_6] = (1-2\gamma) X_4$	
	$[X_5, X_6] = \gamma X_5$		
$\mathfrak{g}_{6,57}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-2\gamma}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = 2\gamma X_2$	$I_2 = \frac{(x_1 x_4 - x_2 x_3)}{x_1^{\gamma+1}}$
	$[X_3, X_6] = (1-\gamma) X_3$	$[X_4, X_6] = \gamma X_4$	
	$[X_5, X_6] = X_4 + \gamma X_5$		
$\mathfrak{g}_{6,58}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2^3 x_1^{-2\gamma}$
	$[X_1, X_6] = 3X_1$	$[X_2, X_6] = 2X_2$	$I_2 = \frac{(x_1 x_4 - x_2 x_3 + \gamma 3x_2^2 \ln(x_1))^3}{x_1^4}$
	$[X_3, X_6] = X_2 + 2X_3$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = \omega X_4 + X_5$		
$\mathfrak{g}_{6,59}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = 0$	$I_2 = x_1 \exp\left(\frac{-x_1 x_4 + x_2 x_3}{x_1 x_2}\right)$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = X_2$	
	$[X_5, X_6] = hX_4$		
$\mathfrak{g}_{6,60}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-2}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = 2X_2$	$I_2 = x_1 \exp\left(\frac{-x_1 x_4 + x_2 x_3}{x_1^2}\right)$
	$[X_3, X_6] = 0$	$[X_4, X_6] = X_1 + X_4$	
	$[X_5, X_6] = \omega X_4 + X_5$		
$\mathfrak{g}_{6,61}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-\lambda}$
	$[X_1, X_6] = 2X_1$	$[X_2, X_6] = 2\lambda X_2$	$I_2 = \frac{x_1 x_4 - x_2 x_3}{x_2 \sqrt{x_1}}$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = (2\lambda - 1) X_4$	
	$[X_5, X_6] = X_3 + X_5$		

Table 10: Nilradical isomorphic to $\mathfrak{g}_{5,1}$ (cont.)

Name	Brackets		Invariants
$\mathfrak{g}_{6,62}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2^2 x_1^{-1}$
$\omega = 0, 1$	$[X_1, X_6] = 2X_1$	$[X_2, X_6] = 2X_2$	$I_2 = x_1 \exp\left(\frac{2(x_1 x_4 - x_2 x_3)}{x_2^2}\right)$
	$[X_3, X_6] = X_2 + X_3$	$[X_4, X_6] = 0$	
	$[X_5, X_6] = \omega X_3 + X_5$		
$\mathfrak{g}_{6,63}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-\lambda}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = \lambda X_2$	$I_2 = x_1 \exp\left(\frac{-x_1 x_4 + x_2 x_3}{x_1 x_2}\right)$
	$[X_3, X_6] = X_3$	$[X_4, X_6] = X_2 + \lambda X_4$	
	$[X_5, X_6] = 0$		
$\mathfrak{g}_{6,64}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 x_1^{-1}$
	$[X_1, X_6] = X_1$	$[X_2, X_6] = X_2$	$I_2 = \frac{(x_2 x_3 - x_1 x_4) + (x_1^2 - h x_2^2) \ln(x_1)}{x_1^2}$
	$[X_3, X_6] = h X_2 + X_3$	$[X_4, X_6] = X_1 + X_4$	
	$[X_5, X_6] = 0$		
$\mathfrak{g}_{6,65}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 \exp^{-\frac{\lambda x_1}{x_2}}$
	$[X_1, X_6] = \lambda X_1 + X_2$	$[X_2, X_6] = \lambda X_2$	$I_2 = (x_1 x_4 - x_2 x_3) \exp^{(\gamma - 2\lambda) x_1 x_2^{-1}}$
	$[X_3, X_6] = (\lambda - \gamma) X_3 + X_4$	$[X_4, X_6] = (\lambda - \gamma) X_4$	
	$[X_5, X_6] = \gamma X_5$		
$\mathfrak{g}_{6,66}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 \exp^{-\frac{2x_1}{x_2}}$
	$[X_1, X_6] = 2X_1 + X_2$	$[X_2, X_6] = 2X_2$	$I_2 = (x_1 x_4 - x_2 x_3) \exp^{-\frac{3x_1}{x_2}}$
	$[X_3, X_6] = X_3 + X_4$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = X_3 + X_5$		
$\mathfrak{g}_{6,67}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 \exp^{-\frac{2x_1}{x_2}}$
	$[X_1, X_6] = 2X_1 + X_2$	$[X_2, X_6] = 2X_2$	$I_2 = (x_1 x_4 - x_2 x_3) \exp^{-\frac{3x_1}{x_2}}$
	$[X_3, X_6] = X_3 + X_4$	$[X_4, X_6] = X_4$	
	$[X_5, X_6] = h X_4 + X_5$		
$\mathfrak{g}_{6,68}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 \exp^{-x_1 x_2^{-1}}$
	$[X_1, X_6] = X_1 + X_2$	$[X_2, X_6] = X_2$	$I_2 = \frac{3(x_1 x_4 - x_2 x_3) x_2 - \gamma x_1^3}{x_2^3}$
	$[X_3, X_6] = X_3 + X_4$	$[X_4, X_6] = c X_1 + X_4$	
	$[X_5, X_6] = 0$		
$\mathfrak{g}_{6,69}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = x_2 \exp^{-x_1 x_2^{-1}}$
	$[X_1, X_6] = X_1 + X_2$	$[X_2, X_6] = X_2$	$I_2 = \frac{2x_1 x_4 - 2x_2 x_3 - x_1^2 - x_2^2}{2} \exp^{-\frac{2x_1}{x_2}}$
	$[X_3, X_6] = X_3 + X_4$	$[X_4, X_6] = X_2 + X_4$	
	$[X_5, X_6] = 0$		
$\mathfrak{g}_{6,70}$	$[X_3, X_5] = X_1$	$[X_4, X_5] = X_2$	$I_1 = (x_1^2 + x_2^2) \left(\frac{i x_1 + x_2}{x_2 - i x_1 l}\right)^{ip}$
	$[X_1, X_6] = p X_1 + X_2$	$[X_2, X_6] = -X_1 + p X_2$	$I_2 = \frac{x_1 x_4 - x_2 x_3}{2} \exp^{(\gamma - 2p) \arctan\left(\frac{x_1}{x_2}\right)}$
	$[X_3, X_6] = (p - \gamma) X_3 + X_4$	$[X_4, X_6] = -X_3 + (p - \gamma) X_4$	
	$[X_5, X_6] = \gamma X_5$		

Table 11: Nilradical isomorphic to $\mathfrak{g}_{5,2}$

Name	Brackets		Invariants
$\mathfrak{g}_{6,71}$	$[X_2, X_5] = X_1$	$[X_3, X_5] = X_2$	$I_1 = \frac{(3x_1^2x_4+x_2^3-3x_1x_2x_3)^{h+1}}{x_1^{h+2}(2x_1x_3-x_2^2)^h}$
	$[X_4, X_5] = X_3$	$[X_1, X_6] = (h+3)X_1$	$I_2 = (-2x_3 + x_2^2x_1^{-1})^{h+3}x_1^{-(h+1)}$
	$[X_2, X_6] = (h+2)X_2$	$[X_3, X_6] = (h+1)X_3$	
	$[X_4, X_6] = hX_4$	$[X_5, X_6] = X_5$	
$\mathfrak{g}_{6,72}$	$[X_2, X_5] = X_1$	$[X_3, X_5] = X_2$	$I_1 = \frac{(3x_1^2x_4+x_2^3-3x_1x_2x_3+x_2^3)^4}{x_1^9}$
	$[X_4, X_5] = X_3$	$[X_1, X_6] = 4X_1$	$I_2 = (-2x_1x_3 + x_2^2)^2x_1^{-3}$
	$[X_2, X_6] = 3X_2$	$[X_3, X_6] = 2X_3$	
	$[X_4, X_6] = X_4$	$[X_5, X_6] = X_4 + X_5$	
$\mathfrak{g}_{6,73}$	$[X_2, X_5] = X_1$	$[X_3, X_5] = X_2$	$I_1 = \frac{3x_1^2x_4+x_2^3-3x_1x_2x_3}{x_1^3}$
$\varepsilon = \pm 1$	$[X_4, X_5] = X_3$	$[X_1, X_6] = X_1$	$I_2 = x_1^\varepsilon \exp\left(\frac{-2x_1x_3+x_2^2}{2x_1^2}\right)$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = \varepsilon X_1 + X_3$	
	$[X_4, X_6] = \varepsilon X_2 + X_4$	$[X_5, X_6] = 0$	
$\mathfrak{g}_{6,74}$	$[X_2, X_5] = X_1$	$[X_3, X_5] = X_2$	$I_1 = \frac{3x_1^2x_4+x_2^3-3x_1x_2x_3}{x_1^3}$
	$[X_4, X_5] = X_3$	$[X_1, X_6] = X_1$	$I_2 = \frac{2x_1x_3-x_2^2}{x_1^2}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = X_3$	
	$[X_4, X_6] = X_4$	$[X_5, X_6] = 0$	
$\mathfrak{g}_{6,75}$	$[X_2, X_5] = X_1$	$[X_3, X_5] = X_2$	$I_1 = x_1^\varepsilon \exp\left(\frac{-3x_1^2x_4-x_2^3+3x_1x_2x_3}{3x_1^3}\right)$
	$[X_4, X_5] = X_3$	$[X_1, X_6] = X_1$	$I_2 = \frac{2x_1x_3-x_2^2}{x_1^2}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = X_3$	
	$[X_4, X_6] = X_1 + X_4$	$[X_5, X_6] = 0$	

Table 12: Nilradical isomorphic to $\mathfrak{g}_{5,3}$

Name	Brackets		Invariants
$\mathfrak{g}_{6,76}$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_3^{(2h+1)} x_1^{-(h+2)}$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = (2h+1) X_1$	$I_2 = \frac{2x_3x_5+x_2^2-2x_4x_1}{x_3x_1^{\frac{h}{2h+1}}} x_1^{2h+1}$
	$[X_2, X_6] = (h+1) X_2$	$[X_3, X_6] = (h+2) X_3$	
	$[X_4, X_6] = X_4$	$[X_5, X_6] = hX_5$	
$\mathfrak{g}_{6,77}$ $\varepsilon = \pm 1$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_1^\varepsilon \exp\left(\frac{-2x_3x_5-x_2^2+2x_1x_4}{2x_1^2}\right)$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = X_1$	$I_2 = x_3x_1^{-2}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = 2X_3$	
	$[X_4, X_6] = \varepsilon X_1 + X_4$	$[X_5, X_6] = 0$	
$\mathfrak{g}_{6,78}$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_1 \exp\left(\frac{2(x_1x_4-x_2x_3)-x_2^2}{2x_1x_3}\right)$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = -X_1$	$I_2 = x_1x_3$
	$[X_2, X_6] = 0$	$[X_3, X_6] = X_3$	
	$[X_4, X_6] = X_3 + X_4$	$[X_5, X_6] = -X_5$	
$\mathfrak{g}_{6,79}$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_1 \exp\left(\frac{-3x_3}{x_1}\right)$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = 3X_1$	$I_2 = \frac{(2(x_1x_4-x_2x_3)-x_2^2)^3}{x_1^4}$
	$[X_2, X_6] = 2X_2$	$[X_3, X_6] = X_1 + 3X_3$	
	$[X_4, X_6] = X_4 + X_5$	$[X_5, X_6] = X_5$	
$\mathfrak{g}_{6,80}$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_3^2x_1^{-1}$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = 2X_1$	$I_2 = \frac{2x_3x_5+x_2^2-2x_1x_4}{x_1}$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = X_3$	
	$[X_4, X_6] = 0$	$[X_5, X_6] = X_5$	
$\mathfrak{g}_{6,81}$ $\varepsilon = \pm 1$	$[X_2, X_4] = X_3$	$[X_2, X_5] = X_1$	$I_1 = x_3^2x_1^{-1}$
	$[X_4, X_5] = X_2$	$[X_1, X_6] = 2X_1$	$I_2 = x_1^\varepsilon \exp\left(\frac{2x_1x_4-2x_2x_3}{x_3^2}\right)$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = X_3$	
	$[X_4, X_6] = 0$	$[X_5, X_6] = \varepsilon X_3 + X_5$	

Table 13: Nilradical isomorphic to $\mathfrak{g}_{5,4}$

Name	Brackets	Invariants
$\mathfrak{g}_{6,82}$ $\alpha = 2$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 2X_1$ $[X_3, X_6] = (\lambda_1 + 1) X_3$ $[X_5, X_6] = (-\lambda_1 + 1) X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = (\lambda + 1) X_2$ $[X_4, X_6] = (-\lambda + 1) X_4$ none
$\mathfrak{g}_{6,82}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = X_1$ $[X_3, X_6] = \lambda_1 X_3$ $[X_5, X_6] = -\lambda_1 X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \lambda X_2$ $[X_4, X_6] = -\lambda X_4$ $I_1 = x_1$ $I_2 = \lambda x_2 x_4 + \lambda_1 x_3 x_5$ $-x_1 x_6$
$\mathfrak{g}_{6,83}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = \left(\frac{\alpha}{2} + \lambda\right) X_3$ $[X_5, X_6] = -X_4 + \left(\frac{\alpha}{2} - \lambda\right) X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \left(\frac{\alpha}{2} + \lambda\right) X_2$ $[X_4, X_6] = \left(\frac{\alpha}{2} - \lambda\right) X_4$ none
$\mathfrak{g}_{6,83}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = \lambda X_3$ $[X_5, X_6] = -X_4 - \lambda X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \lambda X_2$ $[X_4, X_6] = -\lambda X_4$ $I_1 = x_1$ $I_2 = \lambda(x_2 x_4 + x_3 x_5)$ $+x_3 x_4 - x_1 x_6$
$\mathfrak{g}_{6,84}$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = 0$ $[X_5, X_6] = X_3$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_2$ $[X_4, X_6] = -X_4$ $I_1 = x_1$ $I_2 = 2x_1 x_6 + x_3^2 - 2x_2 x_4$
$\mathfrak{g}_{6,85}$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 2X_1$ $[X_3, X_6] = X_3$ $[X_5, X_6] = X_3 + X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = (1 + \lambda) X_2$ $[X_4, X_6] = (1 - \lambda) X_4$ none
$\mathfrak{g}_{6,86}$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 2X_1$ $[X_3, X_6] = X_3$ $[X_5, X_6] = -X_4 + X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_2 + X_3$ $[X_4, X_6] = X_4$ none
$\mathfrak{g}_{6,87}$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 2X_1$ $[X_3, X_6] = X_3 + X_4$ $[X_5, X_6] = X_3 + X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_2 + X_5$ $[X_4, X_6] = X_4$ none
$\mathfrak{g}_{6,88}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = -v_0 X_2 + \left(\frac{\alpha}{2} + \mu_0\right) X_3$ $[X_5, X_6] = -v_0 X_4 + \left(\frac{\alpha}{2} - \mu_0\right) X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \left(\frac{\alpha}{2} + \mu_0\right) X_2 + v_0 X_3$ $[X_4, X_6] = \left(\frac{\alpha}{2} - \mu_0\right) X_4 + v_0 X_5$ none
$\mathfrak{g}_{6,88}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = -v_0 X_2 + \mu_0 X_3$ $[X_5, X_6] = -v_0 X_4 - \mu_0 X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \mu_0 X_2 + v_0 X_3$ $[X_4, X_6] = -\mu_0 X_4 + v_0 X_5$ $I_1 = x_1$ $I_2 = \mu_0(x_2 x_4 + x_3 x_5)$ $+v_0(x_3 x_4 - x_2 x_5) - x_1 x_6$

Table 14: Nilradical isomorphic to $\mathfrak{g}_{5,4}$ (cont.)

Name	Brackets	Invariants
$\mathfrak{g}_{6,89}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = v_0 X_5 + \frac{\alpha}{2} X_3$ $[X_5, X_6] = -v_0 X_3 + \frac{\alpha}{2} X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \left(\frac{\alpha}{2} + s\right) X_2$ $[X_4, X_6] = \left(\frac{\alpha}{2} - s\right) X_4$ none
$\mathfrak{g}_{6,89}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = v_0 X_5$ $[X_5, X_6] = -v_0 X_3$	$[X_3, X_5] = X_1$ $[X_2, X_6] = s X_2$ $[X_4, X_6] = -s X_4$ $I_1 = x_1$ $I_2 = \frac{2x_1 x_6 - 2s x_2 x_4 - v_0(x_3^2 + x_5^2)}{2}$
$\mathfrak{g}_{6,90}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = \frac{\alpha}{2} X_3 + v_0 X_5$ $[X_5, X_6] = -v_0 X_3 + \frac{\alpha}{2} X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \frac{\alpha}{2} X_2 + X_4$ $[X_4, X_6] = X_2 + \frac{\alpha}{2} X_4$ none
$\mathfrak{g}_{6,90}$ $\alpha = 0$ $v_0 \neq 1$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = v_0 X_5$ $[X_5, X_6] = -v_0 X_3$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_4$ $[X_4, X_6] = X_2$ $I_1 = x_1$ $I_2 = \frac{2x_1 x_6 + x_2^2 - x_4^2 - v_0(x_3^2 + x_5^2)}{2}$
$\mathfrak{g}_{6,91}$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = X_5$ $[X_5, X_6] = -X_3$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_4$ $[X_4, X_6] = X_2$ $I_1 = x_1$ $I_2 = \frac{2x_1 x_6 + x_2^2 - x_4^2 - (x_3^2 + x_5^2)}{2}$
$\mathfrak{g}_{6,92}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = -\mu_0 X_2 + \frac{\alpha}{2} X_3$ $[X_5, X_6] = -v_0 X_4 + \frac{\alpha}{2} X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \frac{\alpha}{2} X_2 + v_0 X_3$ $[X_4, X_6] = \mu_0 X_5 + \frac{\alpha}{2} X_4$ none
$\mathfrak{g}_{6,92}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = -\mu_0 X_2$ $[X_5, X_6] = -v_0 X_4$	$[X_3, X_5] = X_1$ $[X_2, X_6] = v_0 X_3$ $[X_4, X_6] = \mu_0 X_5$ $I_1 = x_1$ $I_2 = -x_1 x_6 - \mu_0 x_2 x_5 + v_0 x_3 x_4$
$\mathfrak{g}_{6,92}^*$ $p \neq 0$	$[X_2, X_4] = X_5$ $[X_1, X_6] = p X_1 + X_3$ $[X_3, X_6] = -X_1 + p X_3$ $[X_5, X_6] = 2p X_5$	$[X_1, X_3] = X_5$ $[X_2, X_6] = p X_2 + X_4$ $[X_4, X_6] = -X_2 + p X_4$ none
$\mathfrak{g}_{6,92}^*$ $p = 0$	$[X_2, X_4] = X_5$ $[X_1, X_6] = X_3$ $[X_3, X_6] = -X_1$ $[X_5, X_6] = 0$	$[X_1, X_3] = X_5$ $[X_2, X_6] = X_4$ $[X_4, X_6] = -X_2$ $I_1 = x_5$ $I_2 = x_1^2 + x_2^2 + x_3^2 + x_4^2 - 2x_5 x_6$
$\mathfrak{g}_{6,93}$ $\alpha \neq 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = \alpha X_1$ $[X_3, X_6] = v_0 X_4 + \frac{\alpha}{2} X_3$ $[X_5, X_6] = -v_0 X_2 + \frac{\alpha}{2} X_5$	$[X_3, X_5] = X_1$ $[X_2, X_6] = \frac{\alpha}{2} X_2 + X_4 + v_0 X_5$ $[X_4, X_6] = X_2 - v_0 X_3 + \frac{\alpha}{2} X_4$ none
$\mathfrak{g}_{6,93}$ $\alpha = 0$	$[X_2, X_4] = X_1$ $[X_1, X_6] = 0$ $[X_3, X_6] = v_0 X_4$ $[X_5, X_6] = -v_0 X_2$	$[X_3, X_5] = X_1$ $[X_2, X_6] = X_4 + v_0 X_5$ $[X_4, X_6] = X_2 - v_0 X_3$ $I_1 = x_1$ $I_2 = \frac{-2x_1 x_6 + 2v_0(x_2 x_3 + x_4 x_5) - x_2^2 + x_4^2}{2}$

Table 15: Nilradical isomorphic to $\mathfrak{g}_{5,5}$

Name	Brackets	Invariants
$\mathfrak{g}_{6,94}$ $\lambda + 2 \neq 0$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = (\lambda + 2) X_1$
	$[X_2, X_6] = (\lambda + 1) X_2$	$[X_3, X_6] = \lambda X_3$
	$[X_4, X_6] = 2X_4$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,94}$ $\lambda + 2 = 0$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = 0$
	$[X_2, X_6] = -X_2$	$[X_3, X_6] = -2X_3$
	$[X_4, X_6] = 2X_4$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,95}$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = 2X_1$
	$[X_2, X_6] = X_2$	$[X_3, X_6] = 0$
	$[X_4, X_6] = X_1 + 2X_4$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,96}$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = 3X_1$
	$[X_2, X_6] = 2X_2$	$[X_3, X_6] = X_3$
	$[X_4, X_6] = X_2 + 2X_4$	$[X_5, X_6] = X_3 + X_5$
$\mathfrak{g}_{6,97}$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = 4X_1$
	$[X_2, X_6] = 3X_2$	$[X_3, X_6] = 2X_3 + X_4$
	$[X_4, X_6] = 2X_4$	$[X_5, X_6] = X_5$
$\mathfrak{g}_{6,98}$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_1, X_6] = X_1$
	$[X_2, X_6] = hX_1 + X_2$	$[X_3, X_6] = X_3$
	$[X_4, X_6] = 0$	$[X_5, X_6] = hX_4$

Table 16: Nilradical isomorphic to $\mathfrak{g}_{5,6}$ (cont.)

Name	Brackets	Invariants
$\mathfrak{g}_{6,99}$	$[X_3, X_4] = X_1$	$[X_2, X_5] = X_1$
	$[X_3, X_5] = X_2$	$[X_4, X_5] = X_3$
	$[X_1, X_6] = 5X_1$	$[X_2, X_6] = 4X_2$
	$[X_3, X_6] = 3X_3$	$[X_4, X_6] = 2X_4$
	$[X_5, X_6] = X_5$	